

TECHNICAL DATA SHEET

Product name	UrtINI
Product type	liquid stinging nettle manure (<i>Urtica dioica</i>)
Classification	Basic substance
Packaging	5L, 20L, 250L, 500L, 1000L
Composition	rich in naturally occurring organic compounds
Action	Insecticide, fungicide, acaricide, biostimulant (booster) action against aphids, mites and fungi.

Mode of usage	<ul style="list-style-type: none"> • To be used in open field and in greenhouses or tunnels • Suitable for all seasons • From sowing stage to harvest (until 7 days before harvest) • Preventive use recommended : if applied before any primary contamination, the higher the efficacy
---------------	---

Application dosage

Crop	Pest	Application timing	Application frequency	Dosage (L/ha)
Oilseed rape, radish	diamond-back moth	In spring and summer until harvest	1-6 applications repeat every 10-12 days	5-10
Potato	aphids	Apply in spring and summer until tuber formation	1-5 applications repeat every 7 to 15 days	5-10
	<i>phytophthorans infestans</i>	Apply in spring and summer until tuber formation	1 to 7 applications repeat every 10 to 21 days	5-10
Orchards (apple, prunes, cherry, currents, peach, walnut)	aphids	Throughout the growing cycle until harvest	1 to 5 applications repeat every 7 to 15 days	5-10
Orchards (apple, prunes, peach, cherry)	alternariose, moniliose, botrytis, rhizopus stolonifer (rotting spots)	Apply in spring and summer until harvest	1 to 6 applications repeat every 10 to 21 days	5-10
Orchards (apple, pear)	codling moth	Apply twice in April and once in May	applications every 15 days during the period of high	5-10

			sensitivity for the codling moth	
Vineyard	red mites	Apply in spring and summer	1 to 6 treatments: 3 treatments before flowering and 3 treatments after flowering	5-10
	mildew	Apply in spring and summer	1 to 6 applications repeat every 10 to 21 days	5-10
Beans	aphids	Apply in spring and summer until harvest	1 to 5 applications repeat every 7 to 15 days	5-10
	red mites	Apply in spring and summer until ripening stage	1 to 6 applications repeat every 10 to 21 days	5-10
Salads, cabbage	aphids	Apply in spring and summer until 9 leaf-stage	1 to 5 applications repeat every 7 to 15 days	5-10
Roses	aphids	Apply in spring and summer	1 to 5 applications repeat every 7 to 15 days	5-10

Additional info

- This product is in accordance with EC 1107/2009 and EC 2017/419.
- All info on efficacy against target pests (insects and fungi) in specific crops and the minimum dosages are specified in the review document from the European Commission SANTE/11809/2016.
- This product is authorized for use in organic farming in accordance with EC 2018/848.
- Product is obtained from the anaerobic fermentation of the aeral parts of the stinging nettle plant (food grade quality).

Instructions

- Do not swallow; wash and dry hands after use.
- Store the product in a dry and cool place away from sunlight.
- Keep away from children, (pet) animals and food and beverages.
- This product is intended for professional and hobby use.